

Life on Mars: Past, present, and future

Chris McKay
NASA Ames Research Center
cmckay@mail.arc.nasa.gov


This talk is a compilation based on the work of many colleagues over many years. My thanks and acknowledgements to all of them

Why is Life on other Worlds Interesting?


- The possibility of a second genesis of life:
 - ⇒ comparative biochemistry
 - ⇒ life is common in the universe (yeah!)

- Information about the early planetary environment
- Relevant to the origin of life on Earth

Why Mars?




Evidence for past liquid water Presence of an atmosphere with CO₂ & N₂ Potential for preservation of evidence of life



Viking:

water frost on Mars

water flowed on the surface



Composition of the Martian Atmosphere

Carbon Dioxide (CO₂) 95.3%

• Nitrogen (N₂) 2.7%

• Argon (Ar) 1.6%


• Water Vapor (H₂O) 0.03% - 0.1%

(saturated in places)

• Oxygen (O_2) 0.13%

Carbon Monoxide (CO) 0.07%

Pressure at the two Viking lander sites (Earth = 1013)




Current Mars Missions

Mars Global Surveyor




Mars Odyssey




Mars Exploration Rovers



Mars Express



Mars Reconnaissance Orbiter



Mars Global Surveyor &


Mars Express

Evidence for water flow on Mars




Nanedi

Lower-Limit of Water Mass Fraction on Mars



Minimum estimated water mass fraction in the top 1 meter of the martian surface from GRS on Mars Odyssey. http://grs.lpl.arizona.edu/

Evidence for Water at Meridiani Planum



'Blueberries' are hemitite concretions

More salt Less salt

Layering consistent with water.




Mars is 1/10 the mass of Earth






No plate tectonics Less gravity No magnetic field

Venus and Mars lack the complete cycle





Thermal
Emission
Spectrometer


indicates a dry cold world.

From the *TES top 25 Science* results:

Unweathered volcanic minerals (pyroxene, feldspar, and minor olivine) dominate the spectral properties of martian dark regions. Conversely, no evidence has been found for weathering products above the TES detection limit. This lack of evidence for chemical weathering of the martian surface indicates a geologic history dominated by a cold, dry climate in which mechanical weathering was the dominant form of erosion.

When Mars was wet it was cold

- Evidence of very low erosion: < 10⁻⁹ m/yr, compare to dry valleys 10⁻⁶ m/yr
- Sporadic distribution of valley features
- Unweathered basaltic surface minerals
- Climate modelers have difficulty getting surface temperatures above 0°C.
- No massive surface carbonates detectable by remote sensing.



Gusev Crater

Large crater lake


Large river

Preserved craters inconsistent with rain

The Dry Valleys

•Largest ice-free region in Antarctica

- •Temperatures:
 - -20°C average
- +10°C maximum
- •1-2 cm equivalent H₂O as snow
- •Pressure well above triple point of H₂O



Lake Vanda and the Onyx River in the Antarctic dry valleys


















Snow-based hydrological cycle





Oldest (probable) fossil on Earth: 3.5 Gyr old


courtesy of J.W. Schopf




Fossils are not enough

- Fossils tell us that there was life on Mars
- But not the nature of that life or its relationship, if any, to life on Earth

Is martian life



ARCHAEA



Fossils are not enough for a forensic investigation.

Possible Sources:

- Viable spores in the soil
- Extant subsurface life



- Organisms preserved in amber or salt
- Organisms preserved in permafrost ←


Permafrost in Siberia: 3.5 Myr old and contains viable bacteria




Beacon Valley, Antarctica: Here there may be 8 Myr old ice; -25°C This ice contains viable bacteria and may be the oldest ice on Earth.









From:Smith & McKay, PSS 2005; data from Acuna et al, 1999; Barlow, 1997, Squyres and Carr, 1986

Limits on long term dormancy

- kT: Thermal decay: ~e^{-ΔE/kT} racemization of amino acids degradation of organic material *not important on Mars*, -70°C
- eV: Radiation from crustal U,Th, K ~0.2rad/yr lethal dose for *Deinococcus radiodurans* in 100 Myr *on Mars hundreds of lethal dose over 3.5 Gyr*
- Its dead, Jim




If we find organic material on Mars (or Europa) how can we tell if it was ever <u>alive?</u>


If its like us then easy, less interesting If its alien then hard, but interesting

How do we recognize alien life?






we'll know it when we see it!



The Lego® Principle

- •Biology is largely built from on a small number of components (Lehninger, 1975):
- -20 L amino acids
- -5 nucleotide bases
- -few D sugars, etc.



The Primordial Biomolecules


The building blocks of Earth life.
Could be different for alien life forms.

The amino acids (in un-ionized form)
OH


Could b

Lehninger, Biochemistry, 1975 page 21

Lysine



Abiotic distributions are smooth Biotic distributions are spiked



McKay 2004 PLoS Biol 2(9)1260-12623











Past

Present

Future??


How much warmer?






from Ramanathan et al. J. Geophys. Res. 1988.

The Greenhouse Gases Blanket





Results





 $+\Delta E =$




Before

After

Energy & Time Requirements for Terraforming Mars

Initial State	Final State	Amount	Energy [J m ⁻²]	Solar Energy ^a [years]	Time [years]
		Surface Warming			
CO ₂ (s) at -125°C	CO ₂ (g) at 15°C	200 kPa; 5.4x10 ⁴ kg m ⁻²	3.7x10 ¹⁰	7.9	
Dirt at -60°C	Dirt at 15°C	~10 m; 2x10 ⁴ kg m ⁻²	1.2x109	0.3	
H ₂ O(s) at -60°C	H ₂ O(l) at 15°C	10 m; 1x10 ⁴ kg m ⁻²	5.5x10 ⁹	1.2	
H ₂ O(s) at -60°C	H ₂ O(g) at 15°C	2 kPa; 5.4x10 ² kg m ⁻²	1.6x10 ⁹	0.33	~
			Total:	10	100
		Deep Warming			
H ₂ O(s) at -60°C	H ₂ O(l) at 15°C	500 m; 5x10 ⁵ kg m ⁻²	2.8x10 ¹¹	56	500
		Making O ₂			X
$CO_2(g) + H_2O$	$CH_2O + O_2(g)$	20 kPa; 5.4x10 ³ kg m ⁻²	8x10 ¹⁰	17	100000

Energy divided by the total solar energy reaching Mars in a year, 4.68x10° J m⁻² yr⁻¹ Adapted from McKay et al., 1991. Nature 352, 489-496.






The Astrobiology Questions

(from the first CAN for Astrobiology from NASA HQ 1997)

- 1. How do habitable worlds form and how do they evolve?
- 2. How did living systems emerge?
- 3. How can other biospheres be recognized?
- 4. How have the Earth and its biosphere influenced each other over time?
- 5. How do rapid changes in the environment affect emergent ecosystem properties and their evolution?
- 6. What is the potential for survival and <u>biological</u> evolution beyond the planet of origin? *NEW*

What I cannot create I do not understand.



Richard P. Feynman written on his office blackboard as he left it for the last time in January 1988

